

The local degrees of freedom of higher dimensional pure Chern-Simons theories

Máximo Bañados^{a,b}, Luis J. Garay^b and Marc Henneaux^{a,c}

^(a) *Centro de Estudios Científicos de Santiago, Casilla 16443, Santiago, Chile*

^(b) *Theoretical Physics Group, The Blackett Laboratory, Imperial College, London SW7 2BZ, UK*

^(c) *Université Libre de Bruxelles, Campus Plaine, C.P. 231, B-1050, Bruxelles, Belgium.*

(27 June 1995)

Abstract

The canonical structure of higher dimensional pure Chern-Simons theories is analysed. It is shown that these theories have generically a non-vanishing number of local degrees of freedom, even though they are obtained by means of a topological construction. This number of local degrees of freedom is computed as a function of the spacetime dimension and the dimension of the gauge group.

04.20.Fy, 11.10.Kk

Three-dimensional pure Chern-Simons theory is well known to possess higher dimensional generalizations. These generalisations are theories in $2n + 1$ dimensions constructed from characteristic classes in $2n + 2$ dimensions in exactly the same way as three-dimensional Chern-Simons theory is built out of the four-dimensional characteristic classes. More precisely, if F^a is the curvature 2-form $F^a = dA^a + \frac{1}{2}f_{bc}^a A^b \wedge A^c$ associated to the gauge field 1-form A^a , where f_{bc}^a are the structure constants of the gauge group, and $g_{a_1 \dots a_{n+1}}$ is a rank $n + 1$, symmetric tensor invariant under the adjoint action of the gauge group, then one defines the Chern-Simons Lagrangian \mathcal{L}_{CS}^{2n+1} through the formula

$$d\mathcal{L}_{CS}^{2n+1} = g_{a_1 \dots a_{n+1}} F^{a_1} \wedge \dots \wedge F^{a_{n+1}}. \quad (1)$$

The three-dimensional case is obtained by taking $n = 1$, which yields $d\mathcal{L}_{CS}^3 = g_{ab} F^a \wedge F^b$, where g_{ab} is an invariant metric on the Lie algebra (necessarily proportional to the Killing metric if the Lie algebra is semisimple).

The Chern-Simons action $I = \int_M \mathcal{L}_{CS}^{2n+1}$ is invariant under standard gauge transformations

$$\delta_\epsilon A_\mu^a = D_\mu \epsilon^a. \quad (2)$$

It is also invariant under spacetime diffeomorphisms, $\delta_\eta A_\mu^a = \mathcal{L}_\eta A_\mu^a$, since \mathcal{L}_{CS}^{2n+1} is a $(2n + 1)$ -form. The spacetime diffeomorphisms can also be represented by

$$\delta_\eta A_\mu^a = \eta^\nu F_{\mu\nu}^a. \quad (3)$$

Indeed, these symmetries differ from the Lie derivative only by a gauge transformation and are often called improved diffeomorphisms [1]. If the only symmetries of the Chern-Simons action are the diffeomorphisms (3) and the gauge transformations (2), then we shall say that there is no accidental gauge symmetry. How this translates into an algebraic condition on $g_{a_1 \dots a_{n+1}}$ will be described precisely below.

Of particular interest are the Chern-Simons theories with gauge group $SO(2n + 1, 1)$ or $SO(2n, 2)$ in $2n + 1$ dimensions because they define gravitational theories [2]. For $n = 1$, one recovers the standard Chern-Simons formulation of Einstein gravity with a cosmological constant [3]. For $n > 1$, one gets the Einstein-Hilbert action supplemented by Lovelock terms [4] with definite coefficients. These gravitational theories admit intriguing black hole solutions [5] generalizing the three-dimensional black holes of Ref. [6].

One of the striking features of Chern-Simons theory in three dimensions is the fact that it has no local degrees of freedom. This is because the equations of motion

$$g_{aa_1 \dots a_n} F^{a_1} \wedge \dots \wedge F^{a_n} = 0 \quad (4)$$

reduce to $F^a = 0$ in the three-dimensional case. Thus, the space of solutions of Chern-Simons theory in three dimensions is the finite-dimensional moduli space of flat connections modulo gauge transformations. [Note that the diffeomorphisms lead to no further quotientizing because they vanish on-shell].

Since the higher dimensional Chern-Simons theories are constructed along the same topological pattern as their three-dimensional analog, one may wonder whether they are also

devoid of local excitations and have only global degrees of freedom. One of the purposes of this letter is to explain why this is not the case. We also count explicitly the number of local degrees of freedom as a function of the dimensions of spacetime and of the gauge group. It turns out that the crucial ingredient that controls the whole analysis is the invariant tensor $g_{a_1 \dots a_{n+1}}$.

We start the discussion with the five dimensional case ($n = 2$) and an N -dimensional abelian group ($G = U(1)^N$). This case already contains all the main points that we want to address and is particularly simple because the invariance condition imposes no restriction on the tensor $g_{a_1 \dots a_{n+1}}$. We shall deal with the general situation of an arbitrary gauge group afterwards.

Assume first that there is only one single abelian field. The equations of motion imply $F \wedge F = 0$, i.e. F has at most rank 2. In the generic case, F has exactly rank 2 (in the space of solutions of $F \wedge F = 0$, the solution $F = 0$ has measure zero). Since F is a closed 2-form, one may bring it locally to the canonical form $F = dx^1 \wedge dx^2$ by a diffeomorphism (Darboux theorem for presymplectic forms of rank 2). Thus, the quotient space of the solutions of the equations of motion modulo the gauge transformations (2) and spacetime diffeomorphisms (3) has locally one and only one solution. This implies that the theory has no local degrees of freedom, in agreement with the findings of Ref. [7].

The case of a single abelian gauge field is, however, a poor representative of what happens in the general situation and, in that sense, is somewhat misleading. The reason is that, in contrast with the three-dimensional Chern-Simons theory, we have also used the diffeomorphisms to prove the absence of local degrees of freedom. Indeed, these diffeomorphisms are needed to bring F to its canonical form. But if there are many abelian fields, then there are many F 's to be brought simultaneously to canonical form and this is not possible with a diffeomorphism. Thus, for many ($N > 1$) abelian fields, one expects the existence of local degrees of freedom unless the invariant tensor g_{abc} happens to have been chosen in some peculiar way that enlarges the number of gauge symmetries of the theory (accidental gauge symmetries).

A typical example of a theory with accidental gauge symmetries is obtained by taking all the mixed components of g_{abc} to vanish, so that the action is just the direct sum of N copies of the action for a single abelian field. The theory is then clearly invariant under diffeomorphisms acting independently on each copy and has no degrees of freedom. But there is no reason to take vanishing mixed components for g_{abc} . If these mixed components differ from zero (and cannot be brought to zero by a change of basis), then the action is not invariant under diffeomorphisms acting independently on each gauge field component A^a , because the invariance of the cross terms requires the diffeomorphism parameters for each copy to be equal, thus gluing all of them together in a single symmetry.

In order to substantiate this discussion and to count precisely the number of local degrees of freedom, it is best to turn to the Hamiltonian analysis [8]. To that end, we shall assume that the spacetime manifold M has the topology $\mathfrak{R} \times \Sigma$, where Σ is a four-dimensional manifold. We then decompose the spacetime gauge field 1-form A^a as $A_\mu^a dx^\mu = A_0^a dt + A_i^a dx^i$ where the coordinate t runs over \mathfrak{R} and the x^i are coordinates on Σ . Although there is no spacetime metric to give any meaning to expressions such as timelike or spacelike, we will call time to the coordinate t and we will say that Σ is a spacelike section as shorthand expressions.

It is easy to see that the Chern-Simons action depends linearly on the time derivative of A_i^a ,

$$I = \int_{\mathfrak{R}} \int_{\Sigma} [l_a^i(A_j^b) \dot{A}_i^a - A_0^a K_a], \quad (5)$$

where K_a is given by

$$K_a = -g_{abc} \epsilon^{ijkl} F_{ij}^b F_{kl}^c. \quad (6)$$

The explicit form of the function $l_a^i(A_j^b)$ appearing in Eq. (5) is not needed here but only its “exterior” derivative in the space of spatial connections, which reads

$$\Omega_{ab}^{ij} \equiv \frac{\delta l_b^j}{\delta A_i^a} - \frac{\delta l_a^i}{\delta A_j^b} = -4\epsilon^{ijkl} g_{abc} F_{kl}^c. \quad (7)$$

The equations of motion obtained by varying the action with respect to A_i^a are given by

$$\Omega_{ab}^{ij} \dot{A}_j^b = \Omega_{ab}^{ij} D_j A_0^b, \quad (8)$$

while the variation of the action with respect to A_0^a yields the constraint $K_a = 0$.

Since the action is linear in the time derivatives of A_i^a , the canonically conjugate momenta p_a^i are subject to the $4N$ primary constraints,

$$\phi_a^i = p_a^i - l_a^i \approx 0. \quad (9)$$

These constraints transform in the coadjoint representation of the Lie algebra because the inhomogeneous terms in the transformation laws of p_a^i and l_a^i cancel out.

It turns out to be more convenient to replace the constraints K_a by the equivalent set

$$G_a = K_a - D_i \phi_a^i. \quad (10)$$

The surface defined by $K_a = 0$, $\phi_a^i = 0$ is equivalent to the surface defined by $G_a = 0$, $\phi_a^i = 0$. The new constraints G_a generate the gauge transformations (2), e.g. $\{A_i^a, \int_{\Sigma} \lambda^b G_b\} = D_i \lambda^a$.

The Hamiltonian action takes the form [8],

$$I = \int_{\mathfrak{R}} \int_{\Sigma} [p_a^i \dot{A}_i^a - A_0^a G_a - u_a^i \phi_a^i]. \quad (11)$$

The Poisson bracket among the constraints is given by

$$\{\phi_a^i, \phi_b^j\} = \Omega_{ab}^{ij}, \quad (12)$$

$$\{\phi_a^i, G_b\} = f_{ab}^c \phi_c^i, \quad (13)$$

$$\{G_a, G_b\} = f_{ab}^c G_c, \quad (14)$$

where f_{ab}^c are the structure constants of the Lie algebra, which vanish in the abelian case that we are considering now. It follows from the constraint algebra that there are no further constraints. The consistency condition $\dot{G}_a = 0$ is automatically fulfilled because G_a is first

class while the other consistency equation $\phi_a^i = \Omega_{ab}^{ij} u_j^b = 0$ will just restrict some of the Lagrange multipliers u_j^b .

Equations (13) and (14) reflect that the constraints G_a are the generators of the gauge transformations and that the constraints ϕ_a^i transform in the coadjoint representation. This means, in particular, that the G_a 's are first class, as mentioned above.

The nature of the constraints ϕ_a^i is determined by the eigenvalues of the matrix Ω_{ab}^{ij} . It turns out that the matrix Ω_{ab}^{ij} is not invertible on the constraint surface $K_a = 0$. Indeed, using some simple combinatorial identities, one can prove that K_a given in Eq. (6) and Ω_{ab}^{ij} satisfy the relation,

$$\Omega_{ab}^{ij} F_{kj}^b = \delta_k^i K_a. \quad (15)$$

This equation shows that, on the constraint surface $K_a = 0$, the matrix Ω_{ab}^{ij} has 4 null eigenvectors $(v_k)_j^b = F_{kj}^b$, ($k = 1 \dots 2n$). The corresponding 4 first class constraints, namely

$$H_i \equiv F_{ij}^a \phi_a^j, \quad (16)$$

generate the spatial diffeomorphisms (3). They satisfy the spatial diffeomorphism algebra, up to gauge transformations. The presence of these constraints is of course not a surprise because the Chern-Simons action is invariant under diffeomorphisms for any choice of the invariant tensor g_{abc} .

One could also expect the presence of another first class constraint, namely, the generator of timelike diffeomorphisms. However, as we shall see below, this symmetry is not independent from the other ones and hence its generator is a combination of the first class constraints G_a and H_i .

We now examine whether the first class constraints G_a and H_i are independent and constitute a complete set. This depends on the properties of the invariant tensor g_{abc} and, for a definite choice of g_{abc} , it also depends on the phase space location of the system. This is due to the fact that the constraint surface of the Chern-Simons theory is stratified into phase space regions where the matrix Ω_{ab}^{ij} has different ranks.

We will say that an invariant tensor g_{abc} is *generic* if and only if it satisfies the following condition: There exist solutions F_{ij}^a of the constraints $K_a = 0$ such that

- (i) the matrix F_{kj}^b (with b, j as row index and k as column index) has maximum rank 4, so that the only solution of $\xi^k F_{kj}^b = 0$ is $\xi^k = 0$ and therefore the 4 null eigenvectors $(v_k)_j^b = F_{kj}^b$, ($k = 1 \dots 4$) are linearly independent;
- (ii) the $(4N) \times (4N)$ matrix Ω_{ab}^{ij} has the maximum rank compatible with (i), namely $4N - 4$; in other words, it has no other null eigenvectors besides $(v_k)_j^b = F_{kj}^b$, ($k = 1 \dots 4$).

We will also say that the solutions F_{ij}^a of the constraints $K_a = 0$ that allow for this condition to be satisfied are *generic*. The reason for this name comes from the following observation. For a given generic g_{abc} , a solution fulfilling both conditions (i) and (ii) will still fulfill them upon small perturbations since maximum rank conditions correspond to inequalities and define open regions. Conversely, a solution not fulfilling conditions (i) or (ii), i.e., located on the surface where lower ranks are achieved (defined by equations

expressing that some non trivial determinants vanish), will fail to remain on that surface upon generic perturbations consistent with the constraints. Non-generic solutions of the constraint equations are also of physical interest but will not be considered here (see Ref. [9] for a more complete analysis).

The genericity condition represents the general case in the sense that it defines an open region in the space of the invariant tensors. Indeed, as we have pointed out, these algebraic conditions enforce inequalities. Therefore, to achieve a lower rank, some extra conditions would have to be fulfilled and this would put g_{abc} on a surface of lower dimensionality in the space of the invariant tensors.

The physical meaning of the above algebraic conditions is straightforward. They simply express that the gauge transformations (2) and the spatial diffeomorphisms (3) are independent and that there are no other first class constraints among the ϕ_a^j besides H_i .

In order to illustrate these points and to show that the genericity condition is not self-contradictory and can be actually fulfilled, let us work out a simple example. Take a non-diagonal g_{abc} of the form:

$$g_{a11} = 0, \quad g_{a'b'1} \equiv g_{a'b'} \quad \text{invertible} \quad (17)$$

where $a', b', \dots = 2, 3 \dots N$. Then, the constraints $K_a = 0$ are solved by taking $F_{ij}^{a'} = 0$ and F_{ij}^1 arbitrary. The matrix Ω_{ab}^{ij} has the tensor product form

$$\Omega_{1a}^{ij} = 0, \quad \Omega_{a'b'}^{ij} = g_{a'b'} \epsilon^{ijkl} F_{kl}^1 \quad (18)$$

and is thus of rank $4(N-1)$ provided that F_{ij}^1 is taken to be invertible. The invertibility of F_{ij}^1 also ensures that the only solution of $\xi^k F_{kj}^b = 0$ is $\xi^k = 0$. Therefore, we can conclude that the invariant tensor g_{abc} given in Eq. (17) is generic. Also, this example shows the stratification of phase space. While the solution that we have discussed (with $\det(F_{ij}^1) \neq 0$) is generic, solutions of the same form but with $\det(F_{ij}^1) = 0$ belong to one of these lower dimensional non-generic phase space regions.

Thus, for generic theories, the only first class constraints are $G_a = 0$ and $H_i = 0$, which shows that the generator of timelike diffeomorphisms is not independent from G_a and H_i . This may be verified explicitly by writing the action of a timelike diffeomorphism parametrised by $\xi^\mu = (\xi^0, 0)$ on A_i^a as, see Eq. (3),

$$\delta_\xi A_i^a = \xi^0 F_{i0}^a. \quad (19)$$

Now, the equations of motion (8) are $\Omega_{ab}^{ij} F_{0j}^b = 0$. Since the only zero eigenvectors of the matrix Ω_{ab}^{ij} are F_{kj}^b , there must exist some ζ^k such that $F_{j0}^b = \zeta^k F_{jk}^b$. Inserting this result in Eq. (19), we obtain

$$\delta_\xi A_i^a = \xi^0 \zeta^k F_{ik}^a, \quad (20)$$

which is an improved spatial diffeomorphism with parameter $\xi^0 \zeta^k$.

We can now count the number of local degrees of freedom in the generic case. We have, $2 \times 4N$ canonical variables (A_i^a, p_a^i) , N first class constraints G_a associated to the gauge invariance, 4 first class constraints H_i associated to the (spatial) diffeomorphism invariance, and $4N - 4$ second class constraints (the remaining ϕ_a^i). Hence, we have

$$\frac{1}{2}[8N - 2(N + 4) - (4N - 4)] = 2N - 2 - N \quad (21)$$

local degrees of freedom. The formula does not apply to $N = 1$ because the spatial diffeomorphisms are not independent in that case, as can be checked directly on the canonical form $F = dx^1 \wedge dx^2$. From (21) we see that, for $N = 2$, there are no degrees of freedom. This happens because one does not use all the diffeomorphism invariance to bring the first F^1 to a canonical form. One may then use the residual diffeomorphism invariance to bring the second field strength F^2 to a canonical form also. However, for $N > 2$, there are degrees of freedom.

The analysis has been performed so far in the abelian case. In the non abelian case, the analysis proceeds similarly, but the invariance condition strongly restricts the possible g_{abc} . So one may fear that there could be a conflict between the invariance condition and the genericity condition. This is not the case and we have verified explicitly that the three-index invariant tensor of $SU(p)$ ($2 < p \leq 6$) is generic. Likewise the gravitational Chern-Simons theories in 5 dimensions are also generic and therefore do carry local degrees of freedom (this was anticipated in quite a different way by Chamseddine who analysed perturbations around a non trivial background [2]).

What has been done in 5 dimensions can be repeated in higher (odd) dimensions. Provided the invariant tensor $g_{a_1 \dots a_{n+1}}$ fulfills a genericity condition that is the straightforward generalization of the one appropriate to 5 dimensions, one finds that the canonical formulation of Chern-Simons theory involves $N + 2n$ first class constraints and $2nN - 2n$ second class constraints in the generic case. The first class constraints generate the gauge symmetries (2) and the spatial diffeomorphisms (3). As in 5 dimensions, the timelike diffeomorphisms can be expressed in terms of the other gauge symmetries. Since there are $2nN$ conjugate pairs, the number of local degrees of freedom is equal to

$$\frac{1}{2}[4nN - 2(N + 2n) - (2nN - 2n)] = nN - n - N, \quad (22)$$

where $N > 1$ and $n > 1$. This expression vanishes only for $n = 2$ and $N = 2$. Again, one may also verify that the genericity condition is not self-contradictory by exhibiting invariant tensors that fulfill it. For instance, one may take a direct generalization of Eq. (17). The complete analysis, where the explicit isolation of the second class constraints is performed and the Dirac bracket is computed, will be reported elsewhere [9].

When the invariant tensor $g_{a_1 \dots a_{n+1}}$ is not generic, Ω_{ab}^{ij} has further zero eigenvalues and thus, there are further gauge symmetries. This implies that the number of degrees of freedom is smaller than in the generic case and may even vanish. As we mentioned above, an extreme example is given by N uncoupled abelian gauge fields, where the extra gauge symmetries are diffeomorphisms acting independently on each individual copy.

To conclude, we have shown that higher dimensional Chern-Simons theories, even though constructed along the same topological pattern as in $2 + 1$ dimensions, do have local degrees of freedom provided that the invariant tensor that enters the action fulfills an appropriate genericity condition. This condition implies that there are no accidental gauge symmetries. The result cannot be anticipated by analysing the case of a single abelian field, which is not representative of the general case.

Acknowledgments. We are grateful to Claudio Teitelboim and Jorge Zanelli for useful conversations. M.B. is partially supported by grants 1930910-93 and 1940203-94 from FONDECYT (Chile), a grant from Fundación Andes (Chile), and by institutional support to the Centro de Estudios Científicos de Santiago provided by SAREC (Sweden) and a group of Chilean private companies (COPEC,CMPC,ENERGIS). L.J.G. is supported by a joint fellowship from the Ministerio de Educación y Ciencia (Spain) and the British Council. This work has been supported in part by research funds from the European Community and by a research grant from F.N.R.S.

REFERENCES

- [1] R. Jackiw, *Phys. Rev. Lett* **41**, 1635 (1978)
- [2] A. H. Chamseddine, *Nucl. Phys.* **346**, 213 (1990); *Phys. Lett.* **B233**, 291 (1989)
- [3] A. Achúcarro, P. Townsend, *Phys. Lett.* **B180**, 89 (1986). E. Witten, *Nucl. Phys.* **B311**, 46 (1998)
- [4] D. Lovelock, *J. Math. Phys.* **12**, 498 (1971)
- [5] M. Bañados, C. Teitelboim, J. Zanelli, *Phys. Rev.* **D49**, 975 (1994)
- [6] M. Bañados, C. Teitelboim, J. Zanelli, *Phys. Rev. Lett.* **69**, 1849 (1992). M. Bañados, M. Henneaux, C. Teitelboim, J. Zanelli, *Phys. Rev.* **D48**, 1506 (1993)
- [7] R. Floreanini, R. Percacci, *Phys. Lett.* **B224**, 291 (1989).
- [8] For a recent treatment on the Dirac method see, M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton University Press, Princeton, 1992).
- [9] M. Bañados, L. J. Garay, M. Henneaux, *The dynamical structure of higher dimensional pure Chern-Simons theories*, in preparation.